22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. Frederick Sachs

DD FORM 1473.84 MAR

D'UNCLASSIFIEDAUNI IMITED | SAME AS RPT.

83 APR edition may be used until exhausted.
All other editions are obsolete.

☐ DTIC USERS

SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED

Unclassified

716/831-3289

22b. TELEPHONE (Include Area Code) | 22c. OFFICE SYMBOL

Final Report 22560-LS.

The original aims of the grant were the study of the effect of cytoskeletal reagents on the activity of stretch activated ion channels.

We found the main difficulty in characterizing the responsiveness was a wide scatter in the dose-response curves. The curves are sigmoidal and we found wide variability between the pressure needed to reach the midpoint oif the curve and also in the maximal slope of the curve (sensitivity). In order to reduce the scatter, we decided that we must begin a program of visualizing the patch so that we could actually measure the stress and strain

rather than having to rely on the applied pressure as the independent variable.

We built a high resolution optical setup in which to do patch clamping. To work at high magnification, we had to redesign the manipulators for minimal drift, redesign the manipulator controller for minimal vibration, and redesign the condenser and illuminator for water immersion, arc-lamp, illumination. We had to build special low clearance tissue culture chambers to permit nearly horizontal approach. We had to design and build hardware and software to acquire data off SVHS video tapes with known time synchronization to pressure stimulii. We wrote software to extract digitized images of the patch and software to curve fit the images to geometric models of the patch so that we knew the patch area, radius of curvature and attachment point to the glass pipette. We also had to design and build a high speed pressure clamp to apply controlled peressure steps or ramps to the patch. We also worked out methods to measure changes in the patch capacitance during stretch. With these tools in place, we measured the channel activity, patch capacitance, membrane stress and membrane strain as a function of time following a step in pressure. The results indicated that the much of the scatter disappears when data is plotted as a function of membrane tensions rather than as a function of applied pressure. We also found that the membrane is rather soft with elasticities of ca. 20dyn/cm indicating that we were not stressing the lipid membrane since that would have produced much higher values. Additionally we found the stress induced increase of specific capacity to be about 0.6uF/cm², and independent of membrane area. This capactiance is characteristic of puge lipids, not protein containing membranes which are more in the range of 1.2 uF/cm². Preliminary experiments have indicated that treatment with cytochalasins do not affect any of the observed parameters, contrary to expectations.

During tenure of the grant we also began a study of the ultrastructure of the patch in order of better understand the distribution of stress and the effectrs of cytoskeletal reagents. We developed an entirely new approach using the high voltage electron microscope. This work also required the development of new apparatus and techniques for handling patch pipettes. We now quick freeze the patch, freeze dry it, and examine the whole mount (in the pipette tip) in the HVEM. We are able to visualize cytoskeleton, membrane and what appear to be intramembrane particles. Using colloidal gold labels we can visualize the location of cell membrane receptors such as acetylcholine receptors and lectin binding sites and


elements of the cytoskeleton such as actin and spectrin.

In order to try and measure the change in dimensions of an SA channel associated with the gating step, we built a laser interferometer which can measure changes in position of much less than an Angstrom. This device has not yet been used on the patch, but ht ebasic idea is to cross correlate the channel openings with the scattered light signal.

We also examined a number of different tissues for stretch activated ion channels and found a whole class of SA channels in astrocytes from the rat. We also found a gadolinium

to be a useful blocker of non-selective cation SA channels.

The grant supported some theoretical work on SA channels and acoustical transduction which has appeared in several review papers.



Bibliography of papers not previously submitted.

•

- 1. Yang, X.C., and F. Sachs. 1987. Stretch activated channels in serveral tissues. <u>Biophysical Journal</u>51:252a-252a.
- 2. Sachs, F. 1988. Mechanical Transduction and Stretch-Activated Ion Channels. <u>Biophysical Journal</u>53:410a (Abstr.).
- 3. Yang, X.C., and F. Sachs. 1988. Stretch-activated (SA) ion channel in Xenopus Occytes: permeation and block. Biophys.J.53:412a (Abstr.)
- 4. Sokabe, M., and F.Sachs. 1988. Imaging patches and simultaneous recording of stretch activated channel activity. Int.Conq.Comp.Physiol.Bioch.815.
- 5. Sachs, F. 1988. Mechanically sensitive ion channels in cultured skeletal muscle cells. <u>Int.Cong.Comp.Physiol.Bioch.</u>235.
- 6. Ding, J.P., C.L. Bowman, M.Sokabe, and F. Sachs. 1989. Mechanical transduction in glial cells: SICS and SACS. <u>Biophys.J.</u>55:244a.
- 7. Jung, F., M.J. Song, and F. Sachs. 1987. Patch clamp anatomy: high voltage electron microscopy of in vivo patches. Biophys.J. 51:517a.
- 8. Jung, F., and F. Sachs. 1987. Electron microscopy of the intact cytoskeleton of stretch human erythrocyte membrane. Biophys.J.51:511a.
- 9. Ding, J.-P., X-C. Yang, C.L.Bowman, and F. Sachs. 1988. A Stretch-Activated Ion Channel in Rat Astrocytes in Primary Cell Culture. Soc.Neurosci.Abstr.14, part2:1056. (Abstr.)
- 10. Gao, X., and F. Sachs. 1989. Improving performance of motorized slides for micromanipulation. <u>J.Neurosci.Methods.</u>28:225-227.
- 11. Yang, X-C., and F. Sachs. 1989. Block of Stretch-Activated Ion Channels in Xenopus Occytes by Gadolinium and Calcium Ions. <u>Science</u>243:1068-1071.
- 12. sachs, F. 1989. Stretch-activated ion channels as acoustical transducers. J. Acoust. Soc. Am. 85:829.
- 13. Ruknudin, A., M.Song, A. Auerbach, and F. Sachs. 1989. The structure of patch clamped membranes in high voltage electron microscopy. Proc.Elec.Mic.Soc.Am. 47:936-937.
- 14. sachs, F., and M.Sokabe. 1989. Stress and strain in patch clamped membranes. Int.Conf.Video.Microscopy 37 (Abstr.)
- 15. Sachs, F. 1989. Ion Channels as Mechanical Transducers. In Cell Shape: Determinants, Regulation and Regulatory Role. W.D. Stein, and F. Bronner, editors. Academic Press, San Diego, NY, Berkeley, Boston.63-92.